



This unedited manuscript has been submitted for publication in the Annals of the NYAS. This paper has not been copyedited.

Nitric Oxide at the cross-road of immunoneuroendocrine interactions

Journal:	<i>Annals of the New York Academy of Sciences</i>
Manuscript ID:	draft
Volume Title:	Neuroimmunomodulation: VIIth International ISNIM Congress
Date Submitted by the Author:	n/a
Complete List of Authors:	Rettori, Valeria; School of Medicine, University of Buenos Aires, CEFYBO Fernandez-Solari, Javier; School of Medicine, University of Buenos Aires, CEFYBO Mohn, Claudia E.; School of Medicine, University of Buenos Aires, CEFYBO Zorrilla Zubilete, Maria A.; School of Medicine, University of Buenos Aires, Neuropharmacology Prestifilippo, Juan Pablo; School of Medicine, University of Buenos Aires, CEFYBO de la Cal, Carolina; School of Medicine University of Buenos Aires, CEFYBO De Laurentiis, Andrea; School of Medicine University of Buenos Aires, CEFYBO
Keywords:	Luteinizing Hormone Releasing Hormone, Oxytocin, Corticosterone, Hypothalamus, Posterior Pituitary, Adrenal Gland, Radiation, Submandibular gland, prenatal stress



1
2
3 **Running Title: Nitric Oxide and Neuroimmunoendocrinology**
4
5

6
7 **Title: Nitric Oxide at the cross-road of immunoneuroendocrine interactions**
8
9

10 **Authors: Valeria Rettori^a, Javier Fernandez-Solari^a, Claudia Mohn^a, María A**
11 **Zorrilla Zubilete^b, Carolina de la Cal^a, Juan Pablo Prestifilippo^a, Andrea De**
12 **Laurentiis^a**
13
14

15
16
17 **Affiliations: Center of Pharmacological and Botanical Studies, School of Medicine,**
18 **University of Buenos Aires, Paraguay 2155 16 floor, Buenos Aires, Argentina^a.**

19 **Department of Neuropharmacology, School of Medicine, University of Buenos**
20 **Aires, Paraguay 2155 15 floor; Buenos Aires, Argentina^b.**
21
22

23
24
25
26 **Corresponding author: Dr. Andrea De Laurentiis, Centro de Estudios**
27 **Farmacológicos y Botánicos, Facultad de Medicina, Paraguay 2155 piso 16, Ciudad**
28 **Autónoma de Buenos Aires, AR 1121ABG, Buenos Aires, Argentina. Phone: 54-11-**
29 **4508-3680 int 112. Fax: 54-11-4508-3680 int 106. andredelaurentiis@yahoo.com.**
30
31
32

33
34
35 **Keywords: luteinizing hormone releasing hormone, oxytocin, corticosterone,**
36 **hypothalamus, posterior pituitary, adrenal gland, radiation, submandibular gland,**
37 **prenatal stress**
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Nitric Oxide (NO) was initially described as a mediator of endothelial relaxation, now its participation is recognized in numerous physiological and pathological processes. It was demonstrated that LPS-stimulated CRH release involves NO production. Furthermore, it has been shown that IL-1, TNF- α , IL-6 and IL-2 can stimulate ACTH release from anterior pituitary via NO. Also, we found that NO released from hypothalamic NOergic neurons in response to NE diffuses to LHRH neurons where activates COX and GC with the consequent increase of PGE₂ and cGMP, respectively, that lead to the exocytosis of LHRH granules. During pathological conditions, such as manganese intoxication, NO production is increased leading to increase in LHRH secretion that can advance puberty. In another study we demonstrated that NO reduces oxytocin, as well as vasopressin secretion from the posterior pituitary suggesting its modulatory role during dehydration. There was found an increase in NOS activity and protein in hippocampus and cerebellum in offspring of rats that were submitted to prenatal stress that correlated with behavioral changes in adults. Also NO participates in signal transduction pathways in peripheral tissue in physiological conditions such as in corticosterone release from the adrenal gland. Pathological conditions, such as tumors of head and neck that are treated with radiation are followed by xerostomy. In a rat model, radiation diminished NOS activity in the submandibular gland followed by inhibition in salivary secretion.

In summary, this review describes the wide participation of NO in the cross-talk between neuroendocrine and neuroimmune systems in physiological and pathological processes.

INTRODUCTION

At the end of the 1980s, it was clearly demonstrated that cells can produce nitric oxide (NO) and that this gaseous molecule is a highly reactive free radical with multiple and complex roles within many biological systems. In the present review our particular aim is to describe the role of NO in the field known as neuroimmunomodulation in which, central nervous system (CNS) activity modulates immune system and in turn, the immune system modulates the activity of the nervous system. In fact, during the past years it has been recognized that NO is a key player in the cross talk between both systems.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The first example of neuroimmunomodulation has been known since 1936, from the pioneer work of Selye, in which a noxious stimulus called stress induced the release of adrenocorticotrophic hormone (ACTH) from the pituitary, which in turn released adrenal cortical steroids.¹ The introduction of bacteria into the body causes the release of toxic soluble products of their cell wall such as lipopolisaccharide (LPS), that induces fever and a concomitant increases in plasma cortisol, as well as the synthesis and release of various cytokines, such as interleukin-1 (IL-1), tumor necrosis factor alpha (TNF- α), IL-6, IL-2, interferon gamma (IFN- γ), between others. These cytokines can release corticotrophin-releasing hormone (CRH) that activates ACTH followed by cortisol release.² The cytokines can be produced not only by immune cells, particularly monocytes and macrophages, but also within the brain by glial elements and neurons.³ We have shown an IL-1 immunoreactive neuronal system with cells bodies in the dorsal preoptic area and anterior hypothalamus, and relatively short axons that could not be traced to the median eminence (ME).⁴ In addition, these substances are synthesized within the pituitary itself.⁵

The research of the last decade indicates that cytokines induce NO production and that NO has a powerfull influence on the secretion of hypothalamic peptides and classic synaptic transmitters, such as catecholamines and gamma aminobutyric acid (GABA); and also can suppress or stimulate the release of pituitary hormones directly.⁶

One of the main physiological effects of NO is due to its binding to heme moiety of guanylate cyclase (GC), by altering its conformation and increasing its activity, causing the conversion of guanosine triphosphate (GTP) to cyclic guanosine monophosphate (cGMP). NO also reacts with several other metalloproteins, such as cytochrome P-450 side chain cleavage, which is essential for steroidogenic reactions.⁷ Another target for NO action is the heme group of the enzyme cyclooxygenase (COX). COX metabolizes free arachidonic acid (AA) to prostaglandins (PG) and thromboxanes. There are two known COX forms; COX-1 is constitutively expressed, whereas COX-2 is strongly induced during inflammation, but it also has been shown to be expressed constitutively in the brain.⁷

NO is synthesized by NO synthase (NOS), an enzyme that converts arginine, in the presence of oxygen and several cofactors, into equimolar quantities of citrulline and NO. There are three variants of the enzyme; two of these are constitutively expressed while the other must be induced. The inducible NOS (iNOS), is formed mainly in immune cells, such as macrophages. LPS combines with its receptors on the surface of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

macrophages and other cells inducing the synthesis of iNOS mRNA. LPS also induces mRNA expression of various cytokines, such as IL-1, IL-6 and TNF- α .⁸

One of the constitutive forms was originally characterized in endothelial cells and was therefore known as endothelial NOS (eNOS), while the other constitutive form, originally characterized in neurons was known as neuronal NOS (nNOS). These two isoforms have been found to be distributed more widely than originally thought. The eNOS is found in the caveoli of endothelial cells and is activated following cholinergic stimulation and the consequent increase of intracellular calcium, producing NO, which diffuses to overlying smooth muscle, and activates GC converting GTP to cGMP, which produces vasodilatation⁹. nNOS is found in the cerebellum and various regions of the cerebral cortex and also in various ganglion cells of the autonomic nervous system. Large number of nNOS-containing neurons were also found in the hypothalamus, particularly in the paraventricular nucleus (PVN) and supraoptic nuclei (SON) with axons projecting to the ME and neural lobe, which also contains large amount of nNOS.¹⁰ Because of this distribution in the hypothalamus in regions that contain peptidergic neurons that control pituitary hormone secretion, we studied the role of NO in the hypothalamic-pituitary axis.

Role of NO in the hypothalamic-pituitary axis during infection

As was previously described, the hypothalamic-pituitary response to infection can be mimicked by the injection of bacterial LPS iv or ip. There is a rapid increase in plasma ACTH and prolactin (PRL) within few minutes accompanied by a rapid inhibition of luteinizing hormone (LH) and thyroid-stimulating hormone (TSH).¹¹ Growth hormone (GH) secretion is stimulated in humans but not in rats.⁴ Also, LPS causes the induction of cytokines synthesis and release from cells of the immune system. It has been found that the first cytokine to be released in the rat and in large quantities is TNF- α , that apparently causes the induction of IL-1 synthesis and release that in turn induces secretion of IL-6.¹² Since the response of the pituitary hormones occurs within few minutes, it is obvious that the secretion of cytokines from immune cells in the periphery cannot be responsible for the immediate alterations in pituitary hormone secretion triggered by LPS. Our research demonstrated that ip injection of LPS induced IL-1 β and iNOS mRNA in the brain, anterior pituitary and pineal glands. The induction of both mRNAs occurred in the meninges, the choroids plexus, the circumventricular organs, such as the subfornical organ and the ME and in the

1
2
3 parvocellular neurons of the PVN and arcuate nucleus (AN), areas of particular interest
4 since they contain the hypothalamic releasing and inhibiting hormone producing
5 neurons and also other neurotransmitters controlled by NO. The greatest induction of
6 iNOS occurred in the anterior lobe of the pituitary.⁸ This massive increase in NO
7 production should further increase the effects of NO to maintain the pattern of
8 hypothalamic hormone secretion already induced by LPS.^{13,14}

9
10
11
12
13
14 The initial response to LPS is mediated in the brain by nNOS. There is no
15 participation of iNOS in this initial response. Indeed, the initial response must be due on
16 LPS receptors in areas where the blood-brain barrier is not present, such as the choroid
17 plexus, ME, organum vasculosum laminae terminalis (OVLT) and other
18 circumventricular organs. LPS induced-input to the hypothalamus occurs at least in part
19 by activation of the locus ceruleus that sends noradrenergic axons to the hypothalamus
20 that synapse on cholinergic interneurons in the PVN and activate CRH release.^{15,16} LPS-
21 stimulated CRH release involves NO production, because it can be blocked by
22 inhibitors of all forms of NOS. Also it has been shown that NO activates COX I leading
23 to generation of prostaglandin E₂ (PGE₂), that activates adenylyl cyclase (AC) and
24 therefore increase cAMP. This cyclic nucleotide activates protein kinase A (PKA)
25 which induces exocytosis of CRH secretory granules into hypophyseal portal vessels,
26 activating ACTH release from the corticotrophs of the anterior pituitary gland.¹⁵ NO
27 activates not only COX, but also lipoxygenase (LOX)^{17,18} and GC, which produces
28 cGMP, that increases intracellular calcium that converts membrane phospholipids into
29 AA, the substrate for COX and LOX, generating PG's and leucotrienes, respectively.
30 Activation of CRH by cytokines can be blocked by cyclosporine, probably by blockade
31 of dephosphorylation of NOS by calcineurin rendering NOS inactive.¹⁹ Furthermore, it
32 has been shown that IL-1, TNF- α , IL-6 and IL-2 can directly stimulate ACTH release
33 from anterior pituitary via NO.^{8,20}

51 ***Role of NO in the control of luteinizing hormone releasing hormone release***

52
53 The control of gonadotropin secretion is extremely complex as revealed by the
54 research since the discovery of LHRH²¹, now commonly called gonadotropin-releasing
55 hormone (GnRH). LHRH controls the release of LH and follicle stimulating (FSH) from
56 the pituitary and also induces mating behavior and penile erection in rats.^{22,23} We
57 demonstrated that intracerebroventricular microinjections of NOS inhibitors inhibited
58 pulsatile LH release and mating behavior which indicates that NO controls the pulsatile
59
60

1
2
3 release of LHRH and mating behavior.^{23,24} In *in vitro* experiments, we showed that NO,
4 released from sodium nitropruside (NP), promoted LHRH release from medial basal
5 hypothalamus (MBH), and that this action was blocked by hemoglobin, a scavenger of
6 NO. NP also increases the release of PGE₂ and LOX products that have been shown to
7 play a role in LHRH release.²⁵ Moreover, inhibitors of COX blocked the release of
8 LHRH induced by norepinephrine, providing further evidence for the role of NO in the
9 control of LHRH release via the activation of COX, as we mentioned above.^{26,27} We
10 postulate that the NO released from the NOergic neurons, near the LHRH neuronal
11 terminals, increases the intracellular free calcium required to activate phospholipase A
12 that converts membrane phospholipids to AA, which then can be converted to PGE₂ via
13 COX. The released PGE₂ activates AC increasing cAMP which activates PKA leading
14 to exocytosis of LHRH secretory granules into the hypophyseal portal vessels for
15 transport to the anterior pituitary gland.²⁸

16
17
18
19
20
21
22
23
24
25
26 Norepinephrine (NE) plays a controlling role in LHRH release by acting on
27 hypothalamic NOergic neurons.²⁶⁻²⁸ We determined that the release of NO from the
28 NOergic neurons has a tonic inhibitory action to decrease the release of NE from
29 hypothalamic explants. Presumably, the NO produced by NOergic neurons diffuses to
30 the terminals of the catecholaminergic neurons, where it acts on GC within the
31 terminals to activate the enzyme and cause the production of cGMP. This cGMP may
32 cause a decrease in intracellular free calcium in the cell, which may block the
33 depolarization of the catecholaminergic terminals, inhibiting NE release from storage
34 vesicles.²⁹ We have shown that NO is increased by NE by α_1 receptor stimulation, since
35 it can be blocked by prazosine, an α_1 receptor blocker.²⁸ The NO diffuses to the
36 noradrenergic terminals and inhibits them, generating a negative feedback to terminate
37 the release of NE and consequently contributing to finalize LHRH surge.²⁹

38
39
40
41
42
43
44
45
46
47
48 The principal excitatory transmitter in the central nervous system is glutamic
49 acid (GA). We demonstrated that the mechanism by which GA stimulates LHRH
50 release involves NE release from noradrenergic neurons, since the alpha receptor
51 blocker, phentolamine, blocked the GA-induced LHRH release.³⁰⁻³² Therefore, GA
52 neurons synapses with the noradrenergic terminal which, in turn, synapses with the
53 NOergic neuron, which then generates NO that diffuses to LHRH terminal to stimulates
54 LHRH release.

55
56
57
58
59
60
Gamma-aminobutyric acid (GABA) plays a dual role in the control of LHRH
release in rats.^{33,34} In female rats it inhibits LHRH secretion by acting on LHRH

1
2
3 neurons in the medial preoptic area and it has a stimulatory effect on LHRH secretion
4 from the arcuate nucleus-median eminence region.³³ NO stimulates GABA release from
5 the hypothalamus of adult male rats, and GABA inhibits LHRH release. We have
6 shown that this inhibition is mediated by NO since the inhibitory effect was prevented
7 by hemoglobin or by NG-monomethyl-L-arginine (NMMA), a competitive inhibitor of
8 NOS. Therefore, NO is involved not only in the stimulation of LHRH release induced
9 by NE, but also in its inhibition by inducing GABA release.³⁵

10
11
12
13
14
15
16 Dopamine (DA) is another catecholamine that, as well as NE, stimulates LHRH
17 release, however, the role of dopamine in pulsatile LHRH release is less clear. There are
18 tuberoinfundibular dopaminergic neurons in the hypothalamus which have been shown
19 to stimulate LHRH release from male rat hypothalami *in vitro*³⁶; however, other studies
20 indicate that at certain concentrations and in different hormonal states such as in the
21 castrate rat, both DA and NE can inhibit LHRH release.³⁷ The release of NO from the
22 NOergic neurons in the hypothalamus has a tonic inhibitory action to decrease the
23 release of DA from the tissue. Whether DA is released prior to each LH pulse to
24 stimulate NO and LHRH release has yet to be determined, but this catecholamine may
25 contribute to release of NO and augment pulsatile LHRH release, since DA receptor
26 blockers can block LH release.³⁸

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The presence of an ultra-short negative feedback that controls the episodic
secretion of LHRH was suggested for the first time by Hyppa *et al.* studying FSH
secretion.³⁹ Morphological and molecular evidences show that LHRH receptors are
expressed in LHRH hypothalamic neurons.⁴⁰ However, there was not much evidence
about the signal transduction pathways triggered by the activation of these receptors *in*
vivo. Krsmanovic *et al.* reported a signal transduction mechanism in GT1-7 neurons,
driven by inositol trisphosphate-induced Ca²⁺ mobilization, responding to LHRH
receptors activation.⁴¹ We demonstrated that LHRH produces differential effects on the
mediators of its own release, in MBH incubated *in vitro*, depending on its concentration
in the inter-synaptic space. LHRH 10⁻¹¹M produced an increase in GABA release but at
10⁻⁷M inhibited GA release, in both cases leading to the inhibition of LHRH release.
Based on our results, we propose the existence of different populations of LHRH
receptors that respond with distinct threshold, activating different signal transduction
pathways. Also, it is possible that LHRH affects its own release by activating an
additional neurotransmitters- independent mechanism. Moreover, it was shown that the
inhibitory effect of buserelin on LHRH release started before changes on

1
2
3 neurotransmitters release were observed.⁴² These findings led us to study the effect of
4 LHRH on cellular messengers involved in LHRH release such as NO and PGE₂. LHRH
5 at 10⁻⁷ and 10⁻¹¹M decreased PGE content, after the incubation of MBH explants during
6 30 min. However, LHRH 10⁻⁷M stimulated NOS activity but at 10⁻¹¹M had no effect.
7 Although NO is a known promoter of LHRH release, the stimulatory effect of the
8 higher concentration of LHRH on NOS activity is not contradictory to the ultra-short
9 negative feedback theory, since, as was described above it was proposed that NO has a
10 dual function inducing LHRH exocytosis during the normal surge, but also, increasing
11 GABA release and decreasing NE and DA release to promote the termination of LHRH
12 pulse.³⁵ Additionally, LHRH 10⁻⁷M but not 10⁻¹¹M increased phosphatidylinositol
13 breakdowns, suggesting that LHRH receptors coupled to PLC are involved in this NO-
14 dependent mechanism. Although the mechanism of episodic release of LHRH is not
15 clear to date, the NO-dependent pathway described could be one of the different
16 existing mechanisms that provide redundancy for this important cycle that controls the
17 physiology of reproduction.
18
19
20
21
22
23
24
25
26
27
28
29

30 These data together with those described above indicate that the NO released
31 from NOergic neurons stimulates LHRH release and initiates the pulse, but also acts
32 back on the norepinephrine and dopaminergic neurons to terminate the pulse. This is
33 coupled with the negative feedforward of GABA released by NO to bring about the
34 termination of the pulse.
35
36
37
38
39

40 ***Participation of NO on the effect of Manganese on LHRH release***

41 Manganese (Mn⁺²) is an essential metal which acts as a cofactor for many
42 enzymes and therefore plays important biological functions.⁴³ Nevertheless, high doses
43 of Mn⁺² can cause reproductive dysfunction.⁴⁴ It was reported that chronic
44 administration of low doses of Mn⁺² to female rats resulted in increased serum levels of
45 puberty-related hormones such as LH, FSH and estradiol, and advanced the time of
46 vaginal opening indicating an advance in puberty.⁴⁵ We demonstrate that manganese
47 chloride (MnCl₂) is capable of stimulating LHRH secretion from MBH and this effect is
48 accompanied by an increase in plasma LH levels. Also, MnCl₂ is capable of stimulating
49 DA secretion and NOS activity in the MBH. Since it was shown that DA can induce
50 NOS/NO⁴⁶, and based on the important role of NO in the control of LHRH²⁵ this
51 suggests that DA/NO activation may mediate the MnCl₂-stimulated secretion of LHRH.
52 In addition, we have also found an action of this metal stimulating the inhibitory
53
54
55
56
57
58
59
60

transmitter, GABA. Furthermore, when GABA-A receptors were blocked by bicuculline the release of LHRH elicited by $MnCl_2$ was doubled. We suggest that the action of $MnCl_2$ to induce GABA secretion may represent a neurotoxic effect to ultimately inhibit LHRH secretion. Also, it was shown that $MnCl_2$ stimulates LHRH release from prepubertal female⁴⁵ and male⁴⁷ rats via the activation of guanylyl cyclase and subsequent stimulation of the cGMP/PKG pathway.⁴⁸ In concordance with that, our recent experiments demonstrate that $MnCl_2$ is capable of stimulating LHRH secretion through the activation of the hypothalamic NO/cGMP/protein kinase G pathway.⁴⁹

The role of NO in control of Posterior Pituitary function

The neural lobe of the pituitary is one of the regions richest in NOS in the rat, with a predominance of the nNOS isoform, suggesting that NO may play a role in controlling the release of neuropeptides and neurotransmitters from the posterior pituitary.^{10,50} NADPH-diaphorase, used as a marker of NOS, was co-localized with vasopressin (VP) and oxytocin (OT) in the hypothalamic nuclei that synthesized these hormones.^{51,52} The synthesis of NO by oxytocinergic or vasopressinergic neurons suggests that it may participate in auto- and/or cross-regulation of OT and VP secretion. Our studies indicate that NO donors reduce OT secretion from the neural pituitary lobe and we postulated that released NO may suppress OT secretion through an ultrashort negative feedback mechanism.⁵³ It has been reported that intracerebroventricular administration of L-NAME enhanced plasma levels of both OT and VP.⁵⁴ Since NOS activity increases following salt loading and dehydration, it has been suggested that this increase may provide a negative feedback to prevent overstimulation of OT and VP release.⁵⁵ In conclusion, NO may intervene in the control of OT response to osmotic stimuli. Also, chronic saline ingestion increases hypothalamic tachykinins concentration in SON.⁵⁶ Tachykinins belong to a family of peptides that include substance P and neurokinin A (NKA). They are contained in hypothalamic neurons and nerve fibers and secretory cells of the posterior and anterior pituitary lobes, suggesting that these peptides may have a physiological role in the control of pituitary function.⁵⁷ Some actions of tachykinins are known to be exerted through NO release.⁵⁸ Although we observed an inhibitory effect of NKA on OT release by activation of NOergic neurons, thus suggesting that NKA may have a dual effect on OT release, decreasing it through NO and increasing OT release by an NO-independent mechanism.⁵⁹ Nevertheless, the net effect of NKA on OT release from posterior pituitary seems to be inhibitory. As we

1
2
3 mentioned above, NO has been linked to the release of several neurotransmitters,
4 including glutamate, acetylcholine, dopamine and GABA. Results on the effect of NO
5 on GABA release are contradictory. Our study showed that endogenous NO has an
6 inhibitory effect on GABAergic activity in posterior pituitary⁵⁹ contrary to the effect
7 that we have reported at hypothalamic level.³⁵ GABAergic terminals in the neural and
8 intermediate lobes participate in the control of OT and VP release^{60,61} raising the
9 possibility that NO modulation of GABAergic activity in the posterior pituitary may be
10 involved in the regulation of the secretory function of this pituitary lobe. However,
11 since NO decreased both, OT and GABA release from the posterior pituitary and
12 GABA was reported to inhibit OT release from nerve terminals of the neural lobe, it is
13 possible that NO might influence the release of OT at the level of the cell bodies and on
14 nerve terminals of the neural lobe by different mechanisms. NO is involved, at least
15 partially, in the reduction of the GABAergic activity induced by NKA. The reduction in
16 GABA release may contribute to the stimulatory effect of NKA on lactotroph function.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Role of NO on the corticosterone release from the adrenal gland

ACTH is the major regulator of steroid secretion and synthesis from the adrenal cortex, inducing an acute secretory response within few minutes and increase in steroid synthesis, including the transcription of steroidogenic genes.⁶³ Many clinical and experimental observations indicate that there could be dissociation between plasma ACTH concentrations and cortisol/corticosterone secretion.⁶⁴ Since the sympathetic nervous system responds earlier than the hypothalamus-pituitary-adrenal axis to stressors, corticosterone release prior to the ACTH action may be due to earlier stimulation of the adrenal cortex via the splanchnic nerves, possibly via NO release. The effect of ACTH on adrenal function may involve NO⁶⁵ since NOS was found expressed in the adrenal gland.⁶⁶ Endothelial and neuronal cells have a close anatomical proximity to steroidogenic cells in the adrenal gland, and these cells release NO. In fact, one study suggested that the production of NO by eNOS or nNOS could be effective in regulating the blood flow and steroid release.⁶⁷ Moreover, NO can also be generated within steroid-producing cells and may be involved in steroidogenesis.⁶⁸

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

On the other hand there are controversial reports on the effects of NO and PGs on the steroidogenic pathway.⁶⁸ All of these studies used dispersed cells, thereby eliminating the normal architecture of the gland. NO has been postulated as an autocrine/paracrine regulator of steroidogenesis in several tissues, with effects on adrenal steroid secretion.^{64,68} We investigated the role of NO and the participation of PGE₂ in corticosterone release stimulated from rat adrenal glands by ACTH *in vitro*.⁶⁹ Our results indicate that NO stimulates corticosterone release from adrenals. We believe that NO leads to a rapid release of stored corticosterone from the adrenals, as the released at 15 min was nearly as great as at 30 min, and corticosterone adrenal content was lower than control, indicating that any *de novo* synthesis of corticosterone occurred during this period. The effect of NO and ACTH on acute release of corticosterone could be mediated by PGE₂, since PGE₂ can release corticosterone and this effect is blocked by indomethacin, a COX inhibitor.⁶⁹

We hypothesized that ACTH activates NOS that increases NO that activates COX that generates PGE₂, which in turn, releases corticosterone. The mechanism by which PGE₂ causes exocytosis remains to be determined.

Effect of radiation on NO production in submandibular gland

The exposure to ionising radiation during therapy for head and neck tumours results in alterations of salivary glands such as sialoadenitis and xerostomia followed by dysphagia, mucositis, rampant dental caries, increased tooth decay, oral infections, oesophagitis and gustatory dysfunction.^{70,71} There has been a continuous effort to understand this salivary gland dysfunction; however, the specific mechanisms that underline such damage remain poorly understood. Salivary glands are composed of highly differentiated almost non cycling cells and differences in radio sensitivity of the various cell types have been observed.⁷² In the submandibular gland (SMG), it has been shown that serous cells are far more radiosensitive and this is in concordance with the observed decrease in salivary flow after irradiation.⁷¹

NOS is widely distributed in different regions of SMG.⁷³ In previous publications of our group we have shown that nNOS is clearly present in nerve terminals and together with iNOS in the apical membrane of the excretory and striated ducts of SMG. Interestingly, NOS was absent in the acinar cells.⁷⁴ Previously reported results from our group indicate that NO has a stimulatory effect on salivary secretion⁷⁵,

1
2
3 therefore we hypothesize that radiation could affect salivary secretion by modulating the
4 NOS activity in the SMG. The inducible NOS expressed in immune cells produces the
5 greatest quantities of NO and many macrophages were seen within the gland.⁷⁵ We
6 observed that iNOS activity was significantly reduced after radiation indicating that the
7 decrease in the levels of NO in SMG could mediate the decreased salivary secretion.⁷⁶
8 Also we found an increase in the PGE content in the SMG that could contribute to the
9 decrease in salivary secretion as we have shown previously.⁷⁷ Furthermore, we found an
10 increase in lipid peroxidation, oxidative stress and an increase in mitochondrial NOS
11 expression in SMG that could lead to a tissue damage observed 180 days after radiation.
12 At 365 days post radiation there were structural changes that could account for
13 irreversible damage such as fibrosis with loss of salivary secretion.
14
15
16
17
18
19
20
21
22
23
24

25 *Effect of prenatal stress on NO production in the CNS of the offspring*

26 The developing CNS is especially vulnerable to stress-induced damages. The
27 effect of stress during the first stages of development induces both, immediate and later
28 alterations, which reflect changes in the neuroendocrine, cognitive and behavioral
29 systems. Stress during pregnancy triggers physiological and behavioral abnormalities in
30 the offspring.⁷⁸
31
32
33
34
35

36 The hippocampus is one of the classic structures associated to cognitive
37 processes. However, lately, the cerebellum has been redefined as a structure with a
38 much wider functionality.⁸⁷ Certain investigations in humans and animal models
39 involve the cerebellum in cognitive functions. We evaluated the effect of stress during
40 pregnancy in the offspring at 30, 60 and 90 postnatal days on the nNOS and iNOS
41 activities and protein contents in the hippocampus and cerebellum. Also we tested the
42 learning and memory capabilities at the same postnatal days.⁸⁰
43
44
45
46
47

48 When protein levels from cerebellum and hippocampus were studied in the
49 offspring of stressed animals, an increase in nNOS was found in both tissues at
50 different ages. At the same time, inducible isoform, iNOS expression did not vary in a
51 significant manner in the analyzed tissues.⁸⁹ Also, we found an increase in the activity
52 in calcium-dependent NOS in cerebellum homogenates in prenatal stressed animals
53 respect to controls.⁸⁰
54
55
56
57
58

59 When spatial memory test was performed it showed that offspring from stressed
60 mothers needed more time to complete the task than control offspring animals. The

1
2
3 differences were highly significant at postnatal day 30 and maintained at day 60 and 90.
4 Also, prenatal stress induced a deficit in the inhibitory avoidance task. As predicted, the
5 latency for the control group increased in a significant manner, which reflects that both,
6 information acquisition and consolidation provided by the context were successful. On
7 the contrary, in the prenatal stressed group, no significant differences were observed
8 when the training latency was compared with the test latency. Also, significant
9 differences were found between control and stressed animals for the test latency values
10 from postnatal day 45. So far, it has been demonstrated that, from a behavioral point of
11 view, prenatal stress induced a decrease in the ability to resolve spatial navigation tasks,
12 along with a learning deficit in the inhibitory avoidance test.⁷⁹

13
14
15
16
17
18
19
20
21 The results indicate that prenatal stress model used is able to induce a decrease
22 in the offspring learning and memory capabilities. These alterations correlate with an
23 increase in the expression and activity of the enzyme NOS in different parts of the brain
24 of stressed rats.
25
26
27

28
29 Therefore, we are able to conclude that the changes in NOS profiles and
30 activities might constitute primary events during the development of the nervous
31 system that may manifest in adulthood as significant changes in the animal behavior.
32
33
34

35 CONCLUSIONS

36
37 Nitric Oxide regulates numerous and diverse physiological processes, including
38 reproduction and neurotransmission. The data presented above indicate that there are
39 many areas in the brain where there is regular periodic physiological release of NO
40 throughout the life span. This occurs in the hippocampus, cerebellum, and, in particular,
41 in the hypothalamus, in which NO controls most of the hypothalamic peptidergic
42 neurons (such as CRH, LHRH, GHRH, somatostatin, OT, and VP) and also activates
43 the release of GABA and inhibits that of NE and DA.
44
45
46
47
48

49
50 Because of the ubiquitous nature of NO, the inappropriate release of this
51 mediator has been linked to pathogenesis of a number of disease states including stress,
52 neurodegenerative disorders, inflammation and septic shock. It is already well known,
53 that stress and infections with release of viral or bacterial products, such as LPS, cause
54 the induction of cytokines, which are released and travel through the bloodstream. LPS
55 and the released cytokines combine with their receptors, induce iNOS expression in
56 neurons, glia and many other immune cells, producing high amount of NO, than would
57 be released in basal conditions. NO itself or combined to other free radicals and other
58
59
60

1
2
3 substances produces toxic effects in cells that bring about cell injury and death. On the
4
5 other hand, there appears to be a temporal and functional relation between de
6
7 hypothalamo-pituitary-adrenal (HPA) axis response and NO formation at every level of
8
9 the axis, suggesting a fundamental role of NO as a key modulator of the HPA to
10
11 maintain homeostasis in these pathological states.
12

13 14 **ACKNOWLEDGEMENTS**

15
16 Most of this work was inspired by Prof. Dr. Samuel M. McCann, who spent the
17
18 last years of his life with us. Our deepest gratitude to our “maestro”.

19
20 Also we thank Ana Ines Casella for her administrative assistance.

21
22 This work was supported by Agencia Nacional de Promoción Científica y
23
24 Tecnológica, Argentina, Grant PICT 14264 and Consejo Nacional de Investigaciones
25
26 Científicas y Técnicas, CONICET, Grant PIP 6149.
27

28 **REFERENCES**

- 29
30 1. Selye, H. 1937. The significance of the adrenals for adaptation. *Science* 85:247-248.
31
32 2. Chowder, I., H. T. Hammel, J. Eiseman, R. M. *et al.* 1996. A comparison of the effects of
33
34 environmental and preoptic heating and pyrogen on plasma cortisol levels. *Am J Physiol*
35
36 210: 606-610.
37
38 3. Breder, C. D., C. A. Dinarello, C. B. Saper. 1988. Interleukin-1 immunoreactive
39
40 innervation of the human hypothalamus. *Science* 240(4850):321.
41
42 4. Rettori V., W. L. Dees, J. K. Hiney, *et al.* 1994. An interleukin-1-alpha-like neuronal
43
44 system in the preoptic-hypothalamic region and its induction by bacterial
45
46 lipopolysaccharide in concentrations which alter pituitary hormone release.
47
48 *Neuroimmunomodulation* 1: 251-258.
49
50 5. Spangelo, B. L., P. C. Isakson & R. M. MacLeod. 1990. Production of interleukin-6 by
51
52 anterior pituitary cells is stimulated by increased intracellular adenosine 3', 5'-
53
54 monophosphate and vasoactive intestinal peptide. *Endocrinology* 127: 403-409.
55
56 6. McCann, S. M., M. Kimura, S. Karanth *et al.* 1998. Role of nitric oxide in the
57
58 neuroendocrine responses to cytokines. *Ann. N. Y. Acad. Sci.* 840:174-84.
59
60 7. Hobbs, A. J., A. Higgs & S. Moncada. 1999. Inhibition of nitric oxide synthase as a
potential therapeutic target. *Annu. Rev. Pharmacol Toxicol.* 39: 191-220.
8. Wong, M. L., V. Rettori, A. al-Shehlee *et al.* 1996. Inducible nitric oxide synthase gene
expression in the brain during systemic inflammation. *Nat. Med.* 2:581-584.

- 1
2
3 9. Förstermann, U., I. Gath, P. Schwarz *et al.* 1995. Isoforms of nitric oxide synthase.
4 Properties, cellular distribution and expressional control. *Biochem. Pharmacol.* 50:1321-
5 1332.
6
7
- 8 10. Bredt, D. S., P. M. Hwang & S. H. Snyder. 1990. Localization of nitric oxide synthase
9 indicating a neural role for nitric oxide. *Nature* 347(6295):768-70.
10
- 11 11. McCann, S. M., J. Antunes-Rodrigues, C. R. Franci *et al.* 2000. Role of the hypothalamic
12 pituitary adrenal axis in the control of the response to stress and infection. *Braz. J. Med.*
13 *Biol. Res.* 33: 1121-1131.
14
15
- 16 12. McCann, S. M., M. Kimura, W. H. Yu *et al.* 2001. Cytokines and pituitary hormone
17 secretion. *Vitam. Horm.* 63: 29-62.
18
- 19 13. Wong, M. L., P. B. Bongiorno, V. Rettori *et al.* 1997. Interleukin (IL) 1 beta, IL-1
20 receptor antagonist, IL-10, and IL-13 gene expression in the central nervous system and
21 anterior pituitary during systemic inflammation : pathophysiological implications. *Proc*
22 *Natl. Acad. Sci.* 94 :227-232.
23
24
- 25 14. Wong, M. L., V. Rettori, A. al-Shekhlee *et al.* 1994. Inducible nitric oxide synthase gene
26 expression in the brain during systemic inflammation. *Nat. Med.* 2: 581-584
27
28
- 29 15. Karanth, S., K. Lyson & S. M. McCann. 1993. Role of nitric oxide in interleukin 2-
30 induced corticotropin-releasing factor release from incubated hypothalami. *Proc. Natl.*
31 *Acad. Sci.* 90: 3383-3387.
32
33
- 34 16. McCann, S. M., J. Antunes-Rodrigues, C.R. Franci *et al.* 2000. Role of the hypothalamic
35 pituitary adrenal axis in the control of the response to stress and infection. *Braz. J. Med.*
36 *Biol. Res.* 33: 1121-1131.
37
38
- 39 17. Karanth, S., K. Lyson, M. C. Aguila *et al.* 1995. Effects of luteinizing-hormone-releasing
40 hormone, alpha-melanocyte-stimulating hormone, naloxone, dexamethasone and
41 indomethacin on interleukin-2-induced corticotropin-releasing factor release.
42 *Neuroimmunomodulation* 2: 166-173.
43
44
- 45 18. Lyson, K. & S. M. McCann. 1992. Involvement of arachidonic acid cascade pathways in
46 interleukin-6-stimulated corticotropin-releasing factor release *in vitro*.
47 *Neuroendocrinology* 55: 708-713.
48
49
- 50 19. Karanth, S., K. Lyson & S. M. McCann. 1994. Cyclosporin A inhibits interleukin-2-
51 induced release of corticotropin-releasing hormone. *Neuroimmunomodulation* 1: 82-85.
52
53
- 54 20. McCann, S. M., S. Karanth, A. Kamat *et al.* 1994. Induction by cytokines of the pattern of
55 pituitary hormone secretion in infection. *Neuroimmunomodulation* 1: 2-13.
56
57
- 58 21. McCann, S. M., S. Taleisnik & H. M. Fridman. 1960. LH-releasing activity in
59 hypothalamic explants. *Proc. Soc. Exp. Biol. Med.* 104: 432-434.
60

22. McCann, S. M. & S. R. Ojeda. 1996. The anterior pituitary and hypothalamus. In Textbook of Endocrine Physiology. JE Griffin and SR Ojeda, Eds. Oxford University Press. Oxford. England: 101-133.
23. Mani, S. K., J. M. Allen, V. Rettori *et al.* 1994. Nitric oxide mediates sexual behavior in female rats by stimulating LHRH release. Proc. Natl. Acad. Sci. 91: 6468-6472.
24. McCann, S. M., M. Kimura, S. Karanth *et al.* 2002. Role of nitric oxide in the neuroendocrine response to cytokines. In: Neuroendocrine-immune Interactions. Gaillard RC (ed.). Front Horm Res 29: 117-129.
25. Rettori, V., N. Belova, W. L. Dees *et al.* 1993. Role of nitric oxide in the control of luteinizing hormone-releasing hormone release in vivo and in vitro. Proc. Natl. Acad. Sci. 90: 10130-10134.
26. Canteros, G., V. Rettori, A. Franchi *et al.* 1995. Ethanol inhibits luteinizing hormone-releasing hormone (LHRH) secretion by blocking the response of LHRH neuronal terminals to nitric oxide. Proc. Natl. Acad. Sci. 92: 3416-3420.
27. Rettori, V., M. Gimeno, K. Lyson *et al.* 1992. Nitric oxide mediates norepinephrine-induced prostaglandin E2 release from the hypothalamus. Proc. Natl. Acad. Sci. 89: 11543-11546.
28. Canteros, G., V. Rettori, A. Genaro *et al.* 1996. Nitric oxide synthase content of hypothalamic explants: increase by norepinephrine and inactivated by NO and cGMP. Proc. Natl. Acad. Sci. 93: 4246-4250.
29. Seilicovich, A., M. Lasaga, M. Befumo *et al.* 1995. Nitric oxide inhibits the release of norepinephrine and dopamine from the medial basal hypothalamus of the rat. Proc. Natl. Acad. Sci. 92: 11299-11302.
30. McCann, S. M. & L. Krulich. 1989. Role of transmitters in control of anterior pituitary hormone release. In: Endocrinology, L. DeGroot ed.; 2nd ed. Philadelphia: W. B. Saunders: 117-130.
31. Rettori V, A. Kamat & S. M. McCann. 1994. Nitric oxide mediates the stimulation of luteinizing-hormone releasing hormone release induced by glutamic acid in vitro. Brain Res. Bull. 33:501-503.
32. Kamat A., W. H. Yu, V. Rettori & S. M. McCann. 1995. Glutamic acid induces luteinizing hormone releasing hormone release via alpha receptors. Brain Res Bull 37: 233-235.
33. McCann, S. M. & V. Rettori. 1986. Gamma amino butyric acid (GABA) controls anterior pituitary hormone secretion. Adv. Biochem. Psychopharmacol. 42: 173-179.
34. Masotto C., G. Wisnieski & A. Negro-Vilar. 1989. Different gamma-aminobutyric acid receptor subtypes are involved in the regulation of opiate-dependent and independent luteinizing hormone-releasing hormone secretion. Endocrinology 125: 548-553.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
35. Seilicovich A., B. H. Duvilanski, D. Pisera *et al.* Nitric oxide inhibits hypothalamic luteinizing hormone-releasing hormone release by releasing gamma-aminobutyric acid. *Proc. Natl. Acad. Sci.* 92: 3421-3424.
36. Ojeda S.R., A. Negro-Vilar & S. M. McCann. 1979. Catecholaminergic modulation of luteinizing hormone-releasing hormone release by median eminence terminals in vitro. *Endocrinology* 104: 617-624.
37. Vijayan E. & S. M. McCann. 1978. Re-evaluation of the role of catecholamines in control of gonadotropin and prolactin release. *Neuroendocrinology* 25: 221-235.
38. Ojeda S. R., P.G. Harms, S. M. McCann. 1974. Effect of blockade of dopaminergic receptors on prolactin and LH release: median eminence and pituitary sites of action. *Endocrinology* 94: 1650-1657.
39. Hyppa M., M. Motta & L. Martini. 1971. 'Ultrashort' feedback control of follicle-stimulating hormone-releasing factor secretion. *Neuroendocrinology* 7: 227-235.
40. Krsmanovic L. Z., S. S. Stojilkovic, L. M. Mertz *et al.* 1993. Expression of gonadotropin-releasing hormone receptors and autocrine regulation of neuropeptide release in immortalized hypothalamic neurons. *Proc. Natl. Acad. Sci. USA* 90: 3908-3912.
41. Krsmanovic, L. Z., N. Mores, C. E. Navarro *et al.* 2001. Regulation of Ca²⁺-sensitive adenylyl cyclase in gonadotropin-releasing hormone neurons. *Mol. Endocrinol.* 15: 429-440.
42. Feleder, C., H. Jarry, S. Leonhardt *et al.* 1996. Evidence to suggest that gonadotropin-releasing hormone inhibits its own secretion by affecting hypothalamic amino acid neurotransmitter release. *Neuroendocrinology* 64: 298-304.
43. Keen, C. L., B. Lönnnerdal & L. S. Hurley. (1984). Manganese. In *Biochemistry of the Essential Ultratrace Elements* (E. Frieden, Eds.) pp. 89-132. Plenum Publishing Co., New York.
44. Grey, L. E. & J. W. Laskey. (1980). Multivariate analysis of the effects of manganese on the reproductive physiology and behavior of the male house mouse. *J. Toxicol. Environ. Health* 6: 861-867.
45. Pine, M., B. Lee, R. Dearth *et al.* 2005. Manganese acts centrally to stimulate luteinizing hormone secretion: A potential influence on female pubertal development. *Toxicol. Sci.* 85: 880-885.
46. Melis, M. R., S. Succu & A. Argiolas. 1996. Dopamine agonists increase nitric oxide production in the paraventricular nucleus of the hypothalamus: correlation with penile erection and yawning. *Eur. J. Neurosci.* 8: 2056-2063.
47. Lee B, Pine M, Johnson L *et al.* 2006. Manganese acts centrally to activate reproductive hormone secretion and pubertal development in male rats. *Reproductive Toxicology* 22: 580-85.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
48. Lee, B., J. K. Hiney, M. D. Pine *et al.* 2007. Manganese stimulates luteinizing hormone releasing hormone secretion in prepubertal female rats: hypothalamic site and mechanism of action. *J. Physiol.* 578 : 765-772.
49. Prestifilippo, J. P., J. Fernández-Solari, C. Mohn *et al.* 2007. Effect of manganese on luteinizing hormone-releasing hormone secretion in adult male rats. *Toxicol. Sci.* 97: 75-80.
50. Sagar, S. M. & D. M. Ferreiro. 1987. NADPH diaphorase activity in the posterior pituitary: relation to neuronal function. *Brain Res.* 400: 348–352.
51. Calka, J. & C. H. Block. 1993. Relationship of vasopressin with NADPH-diaphorase in the hypothalamo-neurohypophyseal system. *Brain Res. Bull.* 32: 207–210.
52. Miyagagua, A., H. Okamura & Y. Ibata. 1994. Coexistence of oxytocin and NADPHdiaphorase in magnocellular neurons of the paraventricular and supraoptic nuclei of the rat hypothalamus. *Neurosci. Lett.* 171: 13–16.
53. Rettori, V., G. Canteros, R. Reynoso *et al.* 1997. Oxytocin stimulates the release of luteinizing hormone–releasing hormone from medial basal hypothalamic explants by releasing nitric oxide. *Proc. Natl. Acad. Sci.* 94: 2741–2744.
54. Kadekaro M, Terrell ML, Liu H. *et al.* 1998. Effects of L-NAME on cerebral, vasopressin, oxytocin, and blood pressure responses in hemorrhaged rats. *Am J Physiol* 274: 1070–1077.
55. Liu, Q. S., Y. Jia & G. Ju. 1997. Nitric oxide inhibits neuronal activity in the supraoptic nucleus of the rat hypothalamic slices. *Brain Res. Bull.* 43: 121-125.
56. Larsen, P. J., D. S. Jessop, S. L. Lightman *et al.* 1993. Preprotachykinin A gene expression in distinct hypothalamic and brain stem regions of the rat is affected by chronic osmotic stimulus: A combined immunohistochemical and in situ hybridization histochemistry study. *Brain Res. Bull* 30: 535-545.
57. Nussdorfer, G. G. & L. K. Malendowicz. 1998. Role of tachykinins in the regulation of the hypothalamo-pituitary-adrenal axis. *Peptides* 19: 949-968.
58. Eutamene, H., V. Theodorou, J. Fioramonti *et al.* 1995. Implication of NK1 and NK2 receptors in rat colonic hypersecretion induced by interleukin 1 beta: role of nitric oxide. *Gastroenterology* 109: 483–489.
59. De Laurentiis, A., D. Pisera, B. Duvilanski, *et al.* 2000. Neurokinin A inhibits oxytocin and GABA release from the posterior pituitary by stimulating nitric oxide synthase. *Brain Res. Bull.* 53: 325–330.
60. Crowley, W. & W. Armstrong. 1992. Neurochemical regulation of oxytocin secretion in lactation. *Endocr. Rev.* 13: 33–65.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
61. Sladek, C. & W. Armstrong. 1987. Gamma-aminobutyric acid antagonists stimulate vasopressin release from organ cultured hypothalamo-neurohypophyseal explants. *Endocrinology* 120: 1576–1580.
62. Pisera, D., A. De Laurentiis, B. Duvilanski B *et al.* 1996. Neurokinin A affects the tubero-hypophyseal GABAergic system. *Neuroreport* 7: 2236–2240.
63. Sewer, M. B. & M. R. Waterman. 2003. ACTH modulation of transcription factors responsible for steroid hydroxylase gene expression in the adrenal cortex. *Microsc. Res. Tech.* 61: 300-307.
64. Bornstein, S. R. & G. P. Chrousos. 1999. Clinical review 104: Adrenocorticotropin (ACTH)- and non-ACTH-mediated regulation of the adrenal cortex: neural and immune inputs. *J. Clin. Endocrinol. Metab.* 84: 1729-1736.
65. Nakayama, T., Y. Izumi, M. Soma *et al.*. 1996. A nitric oxide synthesis inhibitor prevents the ACTH-stimulated production of aldosterone in rat adrenal gland. *Endocrin. J.* 43: 157-162.
66. Cymeryng, C. B., S. P. Lotito, C. Colonna, *et al.* 2002. Expression of nitric oxide synthases in rat adrenal zona fasciculata cells. *Endocrinology.* 143: 1235-1242.
67. Riquelme, R. A., G. Sánchez, L. Liberona *et al.* 2002. Nitric oxide plays a role in the regulation of adrenal blood flow and adrenocorticomedullary functions in the llama fetus. *J. Physiol.* 544: 267-276.
68. Cymeryng, C. B., L. A. Dada, C. Colonia *et al.* 1999. Effects of L-arginine in rat adrenal cells: involvement of nitric oxide synthase *Endocrinology* 140: 2962-2967.
69. Mohn, C. E., J. Fernandez-Solari, A. De Laurentiis *et al.* 2005. The rapid release of corticosterone from the adrenal induced by ACTH is mediated by nitric oxide acting by prostaglandin E2. *Proc. Natl. Acad. Sci.* 102: 6213-6218.
70. Valdez, I. H., J. C. Atkinson, J. A. Ship *et al.* 1993. Major salivary gland function in patients with radiation-induced xerostomia: flow rates and sialochemistry. *Int J. Radiat. Oncol. Biol. Phys.* 25: 41-47.
71. Nagler, R. M., B. J. Baum, G. Miller *et al.* 1998. Long term salivary effects of single-dose head and neck irradiation in the rat. *Archs. Oral Biol.* 43: 297-303.
72. O'Connell, A., R. Redman, L. Evans *et al.* 1999. Radiation-induced progressive decrease in fluid secretion in rat submandibular gland is related to decrease acinar volume and not impaired calcium signaling. *Rad. Res.* 151: 150-158.
73. Lohinai, Z., A. D. Szekely, L. Soos *et al.* 1995. Distribution of nitric oxide synthase containing elements in the feline submandibular gland. *Neurosci. Lett.* 192: 9–12.
74. Lomniczi, A., A. M. Suburo, J. C. Elverdin *et al.* 1998. Role of nitric oxide in salivary secretion. *Neuroimmunomodulation* 5: 226–233.


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
75. Rettori, V., A. Lomniczi, J. C. Elverdin *et al.* 2000. Control of salivary secretion by nitric oxide and its role in neuroimmunomodulation. *Ann. N. Y. Acad. Sci.* 917: 258-267.
 76. de la Cal, C., A. Lomniczi, C. E. Mohn *et al.* 2006. Decrease in salivary secretion by radiation mediated by nitric oxide and prostaglandins. *Neuroimmunomodulation* 13: 19-27.
 77. Lomniczi, A., C. Mohn, A. Faletti *et al.* 2001. Inhibition of salivary secretion by lipopolisaccharide : possible role of prostaglandins. *Am. J. Physiol. Endocrinol. Metab.* 281: E405-E411.
 78. Ruiz, R. & C. Avant. 2005. Effects of Maternal Prenatal Stress on Infant Outcomes. *Adv. Nurs. Sci.* 28: 345-355.
 79. Martin, L., D. Goldowitz & G. Mittleman. 2003. The cerebellum and spatial ability: dissection of motor and cognitive components with a mouse model system. *Eur J Neurosci* 18: 2002-2010.
 80. Romero, C. G., D. G. Maur, M. L Palumbo *et al.* 2008. Alteration induced by prenatal stress on NO system in offspring CNS. *Neurochem. Inter.* (in press).

FIGURE LEGEND

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Diagram of the postulated participation of NO in LHRH release. Glutamic acid (Glut) released from its neurons binds to NMDA receptors located on catecholaminergic neurons inducing NE release, which acts on α -adrenergic receptors activating nNOS and therefore the production of NO. NO exerts its stimulatory actions on LHRH release by the interaction with the heme group of, at least, two enzymes. One is the activation of COX, present in LHRH neurons, producing the conversion of arachidonic acid (AA) into PGE₂ that stimulates AC activity with the consequent increase in cAMP production that induces LHRH secretion via cAMP-dependent protein kinase (PKA). The other is activation of GC, increasing cGMP production that stimulates LHRH secretion via cGMP-dependent PKG. Also, by releasing AA from membrane phospholipids, enhances PGE₂ production. On the other hand, NO can produce inhibitory actions on LHRH release. NO depolarizes GABAergic neurons and therefore increases GABA release that acts on GABA receptors located on LHRH neurons producing the inhibition of LHRH release. Also, NO blocks the depolarization of the catecholaminergic terminals, inhibiting NE release and therefore the NE-stimulatory pathway. Both inhibitory actions of NO probably are necessary to terminate LHRH pulses. Solid arrows indicate stimulation. Dashed arrows indicate inhibition.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



210x297mm (200 x 200 DPI)